




## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>C12N 15/82, 9/10, 9/24, 9/88, 9/14, 15/54, 15/55, 15/60, 5/10, A01H 5/00</b>		<b>A1</b>	(11) International Publication Number: <b>WO 99/07863</b>
			(43) International Publication Date: 18 February 1999 (18.02.99)
(21) International Application Number: <b>PCT/EP98/04878</b>		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 5 August 1998 (05.08.98)			
(30) Priority Data: 197 34 213.3      7 August 1997 (07.08.97)      DE			
(71) Applicant (for all designated States except US): MAX-PLANCK-GESELLSCHAFT ZUR FÖRDERUNG DER WISSENSCHAFTEN E.V. [DE/DE]; Berlin (DE).			
(72) Inventors; and (75) Inventors/Applicants (for US only): KWART, Marion [DE/DE]; Ellwanger Strasse 7, D-12247 Berlin (DE). RIESMEIER, Jörg [DE/DE]; Ludwigsfelder Strasse 18, D-14165 Berlin (DE). WILLMITZER, Lothar [DE/DE]; Arnold-Knoblauch-Ring 1, D-14109 Berlin (DE).			
(74) Agent: VOSSIUS & PARTNER; P.O. Box 86 07 67, D-81634 München (DE).		<b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	

(54) Title: PROCESSES FOR INCREASING THE YIELD IN PLANTS



## (57) Abstract

Described are processes for increasing the yield in plants in which nucleotide sequences are expressed under the control of companion cell specific promoters. The nucleotide sequences encode proteins the expression of which leads to a stimulation of the loading of the phloem with photoassimilates.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LJ	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

## Processes for increasing the yield in plants

### Description

The present invention relates to processes for increasing the yield in plants, recombinant nucleic acid molecules used for these processes, their uses as well as to plants with an increased yield.

In the field of agriculture and forestry constantly efforts are being made to produce plants with an increased yield, in particular in order to guarantee the supply of the constantly increasing world population with food and to guarantee the supply of reproducible raw materials. Conventionally, it is tried to obtain plants with an increased yield by breeding, which is, however time-consuming and labor-intensive. Furthermore, appropriate breeding programs have to be performed for each relevant plant species.

Progress has partly been made by the genetic manipulation of plants, that is by introducing into and expressing recombinant nucleic acid molecules in plants. Such approaches have the advantage of usually not being limited to one plant species but being transferable to other plant species. In EP-A 0 511 979, e.g., it was described that the expression of a prokaryotic asparagine synthetase in plant cells *inter alia* leads to an increased biomass production.

In WO 96/21737, e.g., the production of plants with an increased yield by the expression of deregulated or unregulated fructose-1,6-bisphosphatase due to the increase of the photosynthesis rate is described.

Nevertheless, there still is a need of generally applicable processes for improving the yield in plants interesting for agriculture or forestry.

Therefore, the problem underlying the present invention is to provide further processes for increasing the yield in plants.

This problem is solved according to the invention by providing the embodiments characterized in the claims.

Therefore, the present invention relates to a process for increasing the yield in plants, characterized in that recombinant DNA molecules stably integrated into the genome of plants are expressed, comprising

- (a) a region allowing the transcription specifically in the companion cells; and operatively linked thereto
- (b) a nucleotide sequence encoding a polypeptide selected from the group consisting of:
  - (i) proteins with an enzymatic activity that cleaves sucrose;
  - (ii) sucrose transporters;
  - (iii) proteins the activity of which leads to the stimulation of the proton gradient located at the plasma membrane of plant cells; and
  - (iv) citrate synthases (E.C. 4.1.3.7).

It was surprisingly found that the expression of the above-described proteins specifically in the phloem of plants leads to a dramatic increase in yield.

The term "increase in yield" preferably relates to an increase of the biomass production, in particular when determined as the fresh weight of the plant.

Such an increase in yield preferably refers to the so-called "sink" organs of the plant, which are the organs that take up the photoassimilates produced during photosynthesis. Particularly preferred are parts of plants which can be harvested, such as seeds, fruits, storage roots, roots, tubers, flowers, buds, shoots, stems or wood. The increase in yield according to the invention is at least 3 % with regard to the biomass in comparison to non-transformed plants of the same genotype when cultivated under the same conditions, preferably at least 10 % and particularly preferred at least 20 %.

The above-described proteins have in common that when they are expressed in the phloem their biological activity leads to an increased loading of the phloem with photoassimilates.

In the context of the present invention photoassimilates are understood to be sugars and/or amino acids.

According to the invention the nucleotide sequence mentioned in (b) can usually encode a plant protein or a bacterial protein or a protein originating from fungi or animal organisms.

In a preferred embodiment the nucleotide sequence encodes a sucrose synthase (E.C. 2.4.1.13), preferably a plant sucrose synthase, in particular from *Solanum tuberosum*, and particularly preferred the type expressed in the tubers of *S. tuberosum*. Such sequences are, for example, described in Salanoubat and Belliard (Gene 60 (1987), 47-56) and are available in the EMBL gene bank under accession number X67125.

In a further preferred embodiment the nucleotide sequence encodes a sucrose phosphorylase (E.C. 2.4.1.7).

Sequences encoding sucrose phosphorylase are, for example, known from WO 96/24679.

In another preferred embodiment the nucleotide sequence encodes an invertase (E.C. 3.2.1.26), preferably an invertase from a microorganism, in particular from a fungus of the genus *Saccharomyces*, preferably from *S. cerevisiae*. Particularly preferred are sequences encoding a cytosolic invertase (Sonnewald et al., Plant J. 1 (1991), 95-106).

According to the invention a sucrose transporter is understood to be a transporter transporting sucrose in plant systems across a membrane. Such a transporter preferably is of plant origin (for example EMBL gene bank accession number G21319). Particularly preferred the sequence described in (b) encodes a sucrose transporter from spinach (*Spinacia oleracea*), in particular with the sequence of the clone SoSUT1, as, e.g., described in Riesmeier et al. (EMBO J. 11 (1992), 4705-4713).

In a further preferred embodiment the protein that stimulates the proton gradient located at the plasma membrane is a proton ATPase.

In this case, the sequence described in (b) preferably encodes a protein from a microorganism, in particular a fungus of the genus *Saccharomyces*, preferably from *S. cerevisiae*.

In a particularly preferred embodiment the sequence encodes the proton ATPase PMA1 from *S. cerevisiae* (Serrano et al., *Nature* 319 (1986), 689-693; EMBL gene bank) or a version of this proton ATPase from *S. cerevisiae* which is truncated at the 3' end, in particular the ATPase  $\Delta$ PAM1 as described in Example 3 of the present invention.

Alternatively, the nucleotide sequence can also encode a proton ATPase from plants, preferably a proton ATPase from *Solanum tuberosum*.

Particularly preferred are sequences encoding the proton ATPase PHA2 from potato (Harms et al, *Plant Mol. Biol.* 26 (1994), 979-988; EMBL gene bank X76535) or a version of this proton ATPase from potato which is truncated at the 3' end, in particular the ATPase  $\Delta$ PHA2 as described in Example 4 of the present invention.

According to the invention the citrate synthase can be any citrate synthase, for example those from bacteria, fungi, animals or plants. DNA sequences encoding citrate synthase are known, for example, from the following organisms: *Bacillus subtilis* (U05256 and U05257), *E. coli* (V01501), *R. prowazekii* (M17149), *P. aeruginosa* (M29728), *A. anitratum* (M33037) (see Schendel et al., *Appl. Environ. Microbiol.* 58 (1992), 335-345 and references cited therein), *Haloferax volcanii* (James et al., *Biochem. Soc. Trans.* 20 (1992), 12), *Arabidopsis thaliana* (Z17455) (Unger et al., *Plant Mol. Biol.* 13 (1989), 411-418), *B. coagulans* (M74818), *C. burnetti* (M36338) (Heinzen et al., *Gene* 109 (1990), 63-69), *M. smegmatis* (X60513), *T. acidophilum* (X55282), *T. thermophila* (D90117), pig (M21197) (Bloxxham et al., *Proc. Natl. Acad. Sci. USA* 78 (1981), 5381-5385), *N. crassa* (M84187) (Ferea et al., *Mol. Gen. Genet.* 242 (1994), 105-110), *S. cerevisiae* (Z11113, Z23259, M14686, M54982, X00782) (Suissa et al., *EMBO J.* 3 (1984), 1773-1781) and potato (EP 95 91 3066.7).

The numbers in brackets are the corresponding accession numbers in the GenEMBL data base.

The nucleotide sequences according to the invention can generally encode any appropriate proteins from any organism, in particular from plants, fungi, bacteria or animals. The sequences preferably encode proteins from plants or fungi. Preferably, the plants are higher plants, in particular starch or oil storing useful plants, for example potato or cereals such as rice, maize, wheat, barley, rye, triticale, oat, millet, etc., as well as spinach, tobacco, sugar beet, soya, cotton etc.

The fungi preferably are of the genus *Saccharomyces*, *Schizosaccharomyces*, *Aspergillus* or *Neurospora*, in particular *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Aspergillus flavus*, *Aspergillus niger* or *Neurospora crassa*.

In a preferred embodiment of the process according to the invention the region mentioned in (a), which guarantees a companion cell specific transcription, is the promoter of the *rolC* gene from *Agrobacterium rhizogenes*.

This promoter is, for example, described in Schmülling et al. (*Plant Cell* (1989), 665-671) and Kühn (Characterization and localization of the sucrose carrier SUT1 in Solanaceae, Doctoral Thesis (1991), Freie Universität Berlin, biology department). Preferably, the region of the promoter is used that has the nucleotide sequence described in Seq ID No. 1.

Apart from the *rolC* promoter mentioned above the person skilled in the art can without further ado use other promoters for a companion cell specific expression. Further companion cell specific promoters are described in the literature, such as the promoter of the sucrose transporter from *Arabidopsis thaliana* (Truernit and Sauer, *Planta* 196 (1995), 564-570).

Furthermore, for different RNAs and proteins their specific occurrence in the companion cells has been described in the literature (see, for example, Foley et al., *Plant Mol. Biol.* 30 (1996), 687-695; DeWitt, *Plant J.* 1 (1991), 121-128; Stadler et al., *Plant Cell* 7 (1995), 1545-1554). Starting from a known protein it is possible for the person skilled in the art without further ado to isolate the cDNA by means of antibodies or by using oligonucleotides derived from the amino acid sequence (cf.,

e.g., Sambrook et al, Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Laboratory Press (1989), Cold Spring Harbor, NY.). Starting from the cDNAs obtained this way it is furthermore possible to screen established genomic libraries from the corresponding organism and to identify genomic fragments. By comparing the nucleotide sequence of the cDNA and of the genomic clone the position of the promoter can be roughly determined. The specificity of the promoter can be verified in a transgenic situation by using chimeric genes consisting of the promoter and indicator genes, such as the  $\beta$ -glucuronidase (cf., e.g., Kertbundit et al., Proc. Natl. Acad. Sci. USA 88 (1991), 5212-5216).

The process according to the invention can in principle be applied to any plant. Therefore, monocotyledonous as well as dicotyledonous plant species are particularly suitable. The process is preferably used with plants that are interesting for agriculture, horticulture and/or forestry.

Examples thereof are vegetable plants such as, for example, cucumber, melon, pumpkin, egg plant, zucchini, tomato, spinach, cabbage species, peas, beans, etc., as well as fruits such as, for example, pears, apples, etc.

Furthermore, oil storing plants are suitable such as, for example, rape, sunflower, soya. In a particularly preferred embodiment starch storing plants are suitable, in particular such as cereals (rice, maize, wheat, rye, oats, triticale, millet, barley), potato, cassava, sweet potato, etc.

The process can also be applied for sucrose storing plants such as, for example, sugar beet and sugar cane, but also for other useful plants such as, for example, cotton, tobacco, types of wood, wine, hops etc.

The invention further relates to recombinant nucleic acid molecules, containing

- (a) a region allowing the transcription specifically in the companion cells of plants; and operatively linked thereto
- (b) a nucleotide sequence encoding a polypeptide, selected from the group consisting of
  - (i) sucrose synthases;
  - (ii) sucrose phosphorylases;
  - (iii) sucrose transporters;



- (iv) proteins the activity of which leads to the stimulation of the proton gradient located at the plasma membrane of plant cells; and
- (v) citrate synthases.

With regard to the preferred embodiments of such molecules, the same applies to the region mentioned in (a) and the nucleotide sequence mentioned in (b) what was already mentioned above in connection with the process of the invention.

The invention also relates to vectors containing nucleic acid molecules of the invention, in particular those which are suitable for the transformation of plant cells as well as for the integration of foreign DNA into the plant genome.

The present invention further relates to plant cells transformed with a nucleic acid molecule of the invention and containing it stably integrated into the genome. These cells differ from naturally occurring plant cells for example in that a nucleic acid molecule of the invention is integrated into the genome of the cell at a location where it does not naturally occur.

The invention further relates to transgenic plants containing plant cells of the invention and, due to the expression of the recombinant nucleic acid molecule integrated into the genome in the companion cells of the plants, showing an increased yield in comparison with corresponding non-transformed plants that were cultivated under the same conditions.

The present invention further relates to propagation material of plants of the invention containing the above-described plant cells of the invention. The term "propagation material" in particular comprises seeds, fruits, tubers, rhizomes, cuttings, calli, cell cultures, etc.

Finally, the present invention relates to the use of recombinant nucleic acid molecules containing a region allowing the transcription specifically in the companion cells of plants and, operatively linked thereto, a nucleotide sequence encoding a polypeptide selected from the group consisting of:

- (i) proteins with sucrose cleaving enzymatic activity;
- (ii) sucrose transporters;
- (iii) proteins the activity of which leads to the stimulation of the proton gradient located at the plasma membrane; and
- (iv) citrate synthases

for the expression in transgenic plants for increasing the yield.

The encoded proteins preferably are the proteins further described above.

Processes for the transformation of monocotyledonous and dicotyledonous plants are known to the person skilled in the art.

For the introduction of DNA into a plant host cell a variety of techniques is available. These techniques comprise the transformation of plant cells with T-DNA using *Agrobacterium tumefaciens* or *Agrobacterium rhizogenes* as transformation means, the fusion of protoplasts, the injection, the electroporation of DNA, the introduction of DNA by means of the biolistic method as well as further possibilities.

For the injection and electroporation of DNA in plant cells the plasmids do not have to fulfill specific requirements. Simple plasmids such as pUC derivatives can be used.

The use of agrobacteria for the transformation of plant cells has extensively been examined and sufficiently disclosed in the specification of EP-A 120 516, in Hoekema (In: The Binary Plant Vector System Offsetdrukkerij Kanters B.V., Alblaserdam (1985), Chapter V), Fraley et al. (Crit. Rev. Plant. Sci. 4, 1-46) and An et al. (EMBO J. 4 (1985), 277-287).

For the transfer of the DNA to the plant cell plant explants can be co-cultivated with *Agrobacterium tumefaciens* or *Agrobacterium rhizogenes*. From the infected plant material (for example leaf explants, segments of stems, roots but also protoplasts or suspension cultivated plant cells) whole plants can be regenerated in a suitable medium which may contain antibiotics or biozides for the selection of transformed cells. The plants obtained that way can then be examined for the presence of the introduced DNA. Other possibilities for the introduction of foreign DNA using the biolistic method or by protoplast transformation are known (cf., e.g., Willmitzer, L., 1993 Transgenic plants. In: Biotechnology, A Multi-Volume Comprehensive Treatise (H.J. Rehm, G. Reed, A. Pühler, P. Stadler, eds.), Vol. 2, 627-659, VCH Weinheim-New York-Basel-Cambridge).

The transformation of dicotyledonous plants via Ti-plasmid-vector systems with the help of *Agrobacterium tumefaciens* is well-established. Recent studies have indicated that also monocotyledonous plants can be transformed by means of vectors based on *Agrobacterium* (Chan et al., *Plant Mol. Biol.* 22 (1993), 491-506; Hiei et al., *Plant J.* 6 (1994), 271-282; Deng et al., *Science in China* 33 (1990), 28-34; Wilmink et al., *Plant Cell Reports* 11 (1992), 76-80; May et al., *Bio/Technology* 13 (1995), 486-492; Conner and Domisse; *Int. J. Plant Sci.* 153 (1992), 550-555; Ritchie et al., *Transgenic Res.* 2 (1993), 252-265).

Alternative systems for the transformation of monocotyledonous plants are the transformation by means of the biolistic method (Wan and Lemaux, *Plant Physiol.* 104 (1994), 37-48; Vasil et al., *Bio/Technology* 11 (1993), 1553-1558; Ritala et al., *Plant Mol. Biol.* 24 (1994), 317-325; Spencer et al., *Theor. Appl. Genet.* 79 (1990), 625-631), the protoplast transformation, the electroporation of partially permeabilized cells, as well as the introduction of DNA by means of glass fibers.

In particular the transformation of maize is described in the literature several times (cf., e.g., WO95/06128, EP 0 513 849; EP 0 465 875; Fromm et al., *Biotechnology* 8 (1990), 833-844; Gordon-Kamm et al., *Plant Cell* 2 (1990), 603-618; Koziel et al., *Biotechnology* 11 (1993), 194-200). In EP 292 435 and in Shillito et al. (*Bio/Technology* 7 (1989), 581) a process is described with the help of which and starting from a mucus-free, soft (friable) maize callus fertile plants can be obtained.

Prioli and Söndahl (*Bio/Technology* 7 (1989), 589) describe the regenerating and obtaining of fertile plants from maize protoplasts of the Cateto maize inbred line Cat 100-1.

The successful transformation of other cereal species has also been described, for example for barley (Wan and Lemaux, see above; Ritala et al., see above) and for wheat (Nehra et al., *Plant J.* 5 (1994), 285-297).

Once the introduced DNA has been integrated into the genome of the plant cell, it usually is stable there and is also contained in the progenies of the originally transformed cell. It usually contains a selection marker which makes the transformed plant cells resistant to a biozide or an antibiotic such as kanamycin, G 418, bleomycin, hygromycin or phosphinotricin and others. Therefore, the individually chosen marker should allow the selection of transformed cells from cells lacking the introduced DNA.

The transformed cells grow within the plant in the usual way (see also McCormick et al., Plant Cell Reports 5 (1986), 81-84). The resulting plants can be cultured normally. Seeds can be obtained from the plants.

Two or more generations should be cultivated to make sure that the phenotypic feature is maintained stably and is transmitted. Seeds should be harvested to make sure that the corresponding phenotype or other properties are maintained.

**Figure 1** schematically shows the construction of the plasmid pBinRoIC-SS.

**Figure 2a** shows the analysis of the sucrose synthase (SS) activity in leaves of transgenic potato plants which had been transformed with the RoIC-SS construct. The enzyme activity was determined according to Zrenner et al. (Plant J. 7 (1995), 97-107). The activity is indicated in  $\mu\text{mol hexose equivalents}/(\text{min} \times \text{g fresh weight})$ .

The columns represent the average values of three samples per genotype. The standard deviation is also indicated.

**Figure 2b** shows the analysis of the tuber yield of transgenic potato plants which had been transformed with the RoIC-SS construct. The columns represent average values of ten to fifteen plants per genotype. The standard deviation is also indicated. The tuber yield is indicated in g per fresh weight.

**Figure 2c** shows the analysis of the tuber starch of transgenic potato plants that had been transformed with the RoIC-SS construct. For this purpose tubers harvested from ten to fifteen plants per genotype were collected and the starch content of the tubers was determined according to Von Schéele et al. (Landw. Vers. Sta. 127 (1937), 67-96).

**Figure 3** schematically shows the construction of the plasmid pBinRoIC-Suc2.

**Figure 4** schematically shows the construction of the plasmid pBinRoIC- $\Delta\text{PMA1}$ .

**Figure 5** schematically shows the cloning strategy of  $\Delta$ PMA1.

Step from A to B:

The  $H^+$ -ATPase  $\Delta$ PMA1, which was truncated at the 3' end, was amplified via PCR with the PMA1 cDNA as the matrix and complementary internal primers (A). The flanking cleavage sites of the PCR product (B) were introduced via the correspondingly synthesized primers.

Step from B to C:

PstI/NotI digestion and cloning of the PCR fragment into the E. coli vector SK- via PstI/NotI cleavage sites (C).

Step from C to D:

BclI/SpeI digestion of the plasmid SK- $\Delta$ PMA1 and cloning of the fragment into the compatible BamHI/XbaI cleavage sites of pBinRoIC (D)

**Figure 6** shows the results of the polymerase chain reaction with specific oligonucleotides indicating the stable integration of  $\Delta$ PMA1 in the genome of transgenic plants which had been obtained by transformation with the rolC- $\Delta$ PMA2 construct. Size of the PCR product = 730 bp; WT = wildtype; M = marker.

**Figure 7** schematically shows the construction of the plasmid pBinRoIC- $\Delta$ PHA2.

**Figure 8** schematically shows the cloning strategy of  $\Delta$ PHA2.

Step from A to B:

The  $H^+$ -ATPase  $\Delta$ PHA2, which was truncated at the 3' end, was amplified via PCR with the PHA2 cDNA as the matrix and complementary internal primers (A). The flanking cleavage sites of the PCR product (B) were introduced via the correspondingly synthesized primers.

Step from B to C:

PstI/EcoRI digestion and cloning of the PCR fragment into the *E. coli* vector SK- via PstI/EcoRI cleavage sites (C).

Step from C to D:

BglII/SpeI digestion of the plasmid SK- $\Delta$ PHA2 and cloning of the fragment into the compatible BamHI/XbaI cleavage sites of pBinRoIC (D)

- Figure 9** shows the results of the polymerase chain reaction with specific oligonucleotides indicating the stable integration of  $\Delta$ PHA2 in the genome of transgenic plants which had been obtained by transformation with the roIC- $\Delta$ PHA2 construct. Size of the PCR product = 758 bp; WT = wildtype; M = marker.
- Figure 10** schematically shows the construction of the plasmid pBinRoIC-SoSUT1.
- Figure 11** schematically shows the construction of the plasmid pBinRoIC-CiSy.
- Figure 12** shows the results of the determination of the sucrose content in parenchymatic samples of tubers of engrafted potato plants enriched with vascular tissue. The genotypes used for engrafting are the lines RoIC-Suc2-#25 (cytosolic invertase) and wildtype *Solanum tuberosum* var. Désirée. The sucrose content was determined according to Stitt et al. (Methods Enzymol. 174 (1989), 518-522). The columns represent the average values of 12 samples per engrafted type. The standard deviation is indicated. The sucrose content is indicated as  $\mu$ mol hexose equivalents/g fresh weight.
- Figure 13** shows the analysis of phloem exudates of  $\Delta$ PMA1 leaves which were kept under light for six hours in a  $^{14}\text{CO}_2$  atmosphere. The sucrose content was determined according to Stitt et al. (loc. cit). The columns represent the average values of four to five samples per genotype. The standard deviation is indicated.

**Figure 14** shows the tuber yield (in gram fresh weight) of  $\Delta$ PMA1 plants. The columns represent the average values of six plants per genotype. The standard deviation is indicated. The tuber yield is indicated in g fresh weight.

**Figure 15** shows the tuber yield (in gram fresh weight) of  $\Delta$ PHA2 plants. The columns represent average values of four to five plants per genotype. The standard deviation is indicated. The tuber yield is indicated in g fresh weight.

The following examples illustrate the invention.

### **Example 1**

#### **Production of the plasmid pBinRoIC-SS and production of transgenic potato plants**

The plasmid pBinRoIC-SS contains the three fragments A, B and C in the binary vector pBin19 (Bevan, Nucl. Acids Res. 12 (1984), 8711) (cf. Fig.1).

The fragment A comprises the roIC promoter from *Agrobacterium rhizogenes*. The roIC promoter contains as an EcoRI/Asp718 DNA fragment of 1138 bp (Lerchl et al., Plant Cell 7 (1995), 259-270) the DNA region (position: 11306 to position 12432) of the TL-DNA of the Ri-agropin-type plasmid from *A. rhizogenes* (Slightom et al., J. Biol. Chem. 261 (1986), 108-121). The fragment A is inserted into the EcoRI and Asp718 cleavage sites of the polylinker of pBin19.

The fragment B contains the coding region (position: 76 to position 2493) of the cDNA of the sucrose synthase (SS) from *Solanum tuberosum* (Salanoubat and Belliard, Gene 60 (1987), 47-56). The fragment B was obtained as BamHI fragment of 2427 bp from the vector pBluescript SK<sup>+</sup>, in which it is inserted into the BamHI cleavage site of the polylinker. The fragment B was inserted in sense orientation in the vector pBin19 into the BamHI cleavage site, that is downstream of the roIC promoter in an orientation allowing the transcription of a translatable RNA.

The fragment C contains the polyadenylation signal of the Gene 3 of the T-DNA of the Ti plasmid pTi ACH 5 (Gielen et al., EMBO J. 3 (1984), 835-846), in particular the nucleotides 11749-11939, which was isolated as a PvuII/HindIII fragment from the plasmid pAGV 40 (Herrera-Estrella et al., Nature 303 (1983), 209-213) and which upon addition of SphI linkers was cloned into the PvuII cleavage site between the SphI and the HindIII cleavage site of the polylinker of pBin19.

The resulting plasmid pBinRoIC-SS was introduced into potato plant cells via the gene transfer mediated by *Agrobacterium tumefaciens*. For this purpose ten small leaves of a potato sterile culture (*Solanum tuberosum* L. cv. Désirée) wounded with the scalpel were put into 10 ml MS medium (Murashige and Skoog, *Physiol. Plant.* 15 (1962)), 473 with 2% of sucrose containing 50 µl of a *Agrobacterium tumefaciens* overnight culture grown under selection. After 3 to 5 minutes of gentle shaking a further incubation followed for two days in the dark. Then the leaves were put on MS medium with 1.6 % glucose, 5 mg/l naphthyl acetic acid, 0.2 mg/l benzylaminopurin, 250 mg/l claforan, 50 mg/l kanamycin and 0.8 % bacto-agar for callus induction. After an incubation of one week at 25 °C and 3000 lux the leaves were put on MS medium with 1.6 % glucose, 1.4 mg/l zeatin ribose, 20 µg/l naphthyl acetic acid, 20 µg/l giberellic acid, 250 mg/l claforan, 50 mg/l kanamycin and 0.8 % bacto-agar.

The analysis of the leaves of a number of plants transformed with this vector system unambiguously indicated the presence of an increased sucrose synthase activity. This is a result of the expression of the sucrose synthase gene from potato contained in pBinRoIC-SS (cf. Figure 2a).

The analysis of the tuber yield (tuber fresh weight in gram) of plants transformed with this vector system and showing an increased sucrose synthase activity unambiguously showed an increased tuber yield. This is also a result of the expression of the sucrose synthase gene from potato contained in pBinRoIC-SS (cf. Figure 2b).

The starch content of potato tubers is linearly dependent on the density of the tubers (von Schéele et al., *Landw. Vers. Sta.* 127 (1937), 67-96). The analysis of the density of transgenic tubers of plants which had been transformed with the vector system pBinRoIC-SS having an increased sucrose synthase activity surprisingly showed an



increased starch content. This is a result of the expression of the sucrose synthase gene from potato contained in pBinRoIC-SS (cf. Figure 2c).

## **Example 2**

### **Production of the plasmid pBinRoIC-Suc2 and production of transgenic potato plants**

The plasmid pBinRoIC-Suc2 contains the three fragments A, B and C in the binary vector pBin19 (Bevan, loc. cit.) and is illustrated in Figure 3.

The fragments A and C correspond to the fragments A and C as described in Example 1.

The fragment B contains the coding region (position: 845 to position: 2384) of the gene of the cytosolic invertase from yeast (*Saccharomyces cerevisiae*). The fragment B was obtained as a BamHI fragment with a length of 1548 bp from the vector pBluescript SK<sup>+</sup> in which it is inserted in the BamHI cleavage site of the polylinker. The fragment B is inserted in sense orientation into pBin19 in the BamHI cleavage site.

The plasmid pBinRoIC-Suc2 was introduced into potato plant cells via gene transfer mediated by *Agrobacterium*. From transformed cells whole plants were regenerated. Such plants show in comparison to non-transformed plants an increased yield (increased biomass).

## **Example 3**

### **Production of the plasmid pBinRoIC- $\Delta$ PMA1 and production of transgenic potato plants**

The plasmid pBinRoIC- $\Delta$ PMA1 contains the three fragments A, B and C in the binary vector pBin19 (Bevan, loc. cit.) and is schematically illustrated in Figure 4.

The fragments A and C correspond to the fragments A and C as described in Example 1.

The fragment B contains the coding region (position: 937 to position: 3666) of the gene of the proton ATPase PMA1 from the yeast *Saccharomyces cerevisiae* (Serrano et al., Nature 319 (1986), 689-693). The fragment B was obtained by means of polymerase chain reaction (PCR). For this purpose the 3' end of the coding region of the gene PMA1 was truncated on purpose by 27 bp and at the same time a necessary new stop codon was introduced. The DNA fragment modified this way was called  $\Delta$ PMA1. The fragment B was inserted, as a BclI/SpeI fragment with a length of 2739 bp, in sense orientation into the BamHI (compatible insertion site for BclI restriction sites) and XbaI (compatible insertion site for SpeI restriction sites) cleavage sites of the vector pBin19.

The fragment B was obtained as BclI/SpeI fragment from the vector pBluescript SK<sup>+</sup> in which it is inserted via the cleavage sites NotI and PstI of the polylinker (cf. Fig. 5). The plasmid pBinRoIC- $\Delta$ PMA1 was introduced into potato plant cells via the gene transfer mediated by *Agrobacterium*. Whole plants were regenerated from transformed cells.

The stable integration of  $\Delta$ PMA1 in the genome of transgenic plants which had been obtained by using the vector system pBinRoIC- $\Delta$ PMA1 was detected by means of polymerase chain reaction (PCR) (cf. Fig. 6).

The transformed plants show an increased yield (increased biomass) in comparison to non-transformed plants (see Figures 13 and 14).

#### **Example 4**

##### **Production of the plasmid pBinRoIC- $\Delta$ PHA2 and production of transgenic potato plants**

The plasmid pBinRoIC- $\Delta$ PHA2 contains the three fragments A, B and C in the binary vector pBin19 (Bevan, loc. cit.) and is schematically illustrated in Figure 7.

The fragments A and C correspond to the fragments A and C as described in Example 1.

The fragment B contains the coding region (position: 64 to position: 2672) of the cDNA of the proton-ATPase PHA2 (Harms et al., Plant Mol. Biol. 26 (1994), 979-988). The fragment B was obtained by means of polymerase chain reaction (PCR). For this purpose the 3' end of the coding region of the gene PHA2 was on purpose truncated by 249 bp, and at the same time two new stop codons were introduced. The DNA fragment modified that way was called  $\Delta$ PHA2. The fragment B was inserted in sense orientation as a BglII/SpeI fragment with a length of 2631 bp into the BamHI (compatible insertion site for BglII restriction sites) and XbaI (compatible insertion site for SpeI restriction sites) cleavage sites of the vector pBin19.

The fragment B was obtained as BglII/SpeI fragment from the vector pBluescript SK<sup>-</sup>, in which it is inserted into the EcoRI and PstI cleavage sites of the polylinker sequence (cf. Fig. 8: cloning strategy  $\Delta$ PHA2).

The plasmid pBinRoIC- $\Delta$ PHA2 was introduced into potato plant cells via the gene transfer mediated by Agrobacterium. Whole plants were regenerated from transformed cells.

The stable integration of  $\Delta$ PHA2 in the genome of transgenic plants which had been obtained using the vector system pBinRoIC- $\Delta$ PHA2 was detected by means of polymerase chain reaction (PCR) (cf. Fig. 9).

The transformed plants show an increased yield (increased biomass) in comparison to non-transformed plants (see Figure 15).

## **Example 5**

### **Production of the plasmid pBinRoIC-SoSUT1 and production of transgenic potato plants**

The plasmid pBinRoIC-SoSUT1 contains the three fragments A, B and C in the binary vector pBin19 (Bevan, loc. cit.) and is schematically illustrated in Figure 10.

The fragments A and C correspond to the fragments A and C as described in Example 1.

The fragment B contains the cDNA (position: 1 to position: 1969) encoding a sucrose transporter from spinach (*Spinacia oleracea*) (Riesmeier et al., EMBO J. 11 (1992), 4705-4713; accession number X67125 and S51273). The fragment B was obtained as a NotI fragment from the vector pBluescript SK<sup>+</sup>, in which it is inserted via a NotI linker sequence. For the cloning into the SmaI cleavage site of the vector pBin19 the sticky ends of the fragment resulting from the NotI digestion were converted to blunt ends and inserted in sense orientation into the pBin19. The resulting plasmid was called pBinRoIC-SoSUT1.

It was introduced into potato plant cells via the gene transfer mediated by *Agrobacterium*. Whole plants were regenerated from transformed cells.

Plants transformed that way show an increased yield (increased biomass) in comparison to non-transformed plants.

### Example 6

#### **Production of the plasmid pBinRoIC-CiSy and production of transgenic potato plants**

The plasmid pBinRoIC-CiSy contains the three fragments A, B and C in the binary vector pBin19 (Bevan, Nucl. Acids Res. 12 (1984), 8711) modified according to Becker (Nucl. Acids Res. 18 (1990), 203) (cf. Fig. 11).

The fragment A comprises the roIC promoter from *Agrobacterium rhizogenes*. The roIC promoter contains as an EcoRI/Asp718 DNA fragment with a length of 1143 bp (Lerchl et al., The Plant Cell 7 (1995), 259-270) the DNA region (position: 11306 to position 12432) of the TL-DNA of the Ri-agropin type plasmid from *A. rhizogenes* (Slightom et al., J. Biol. Chem. 261 (1986), 108-121). The fragment A is inserted in the EcoRI and Asp718 cleavage sites of the polylinker of pBin19.

The fragment B contains the coding region of the cDNA of the citrate synthase (CiSy) from the fission yeast *Saccharomyces cerevisiae*. The fragment B was obtained as a BamHI fragment with a length of 1400 bp from the vector pBluescript SK<sup>-</sup>, in which it is inserted in the BamHI cleavage site of the polylinker (Landschütze, Studies on the influence of the acetyl-CoA synthesis and use in transgenic plants, Doctoral Thesis, Freie Universität Berlin, (1985) D83/FB15 No. 028).

The fragment C contains the polyadenylation signal of the gene 3 of the T-DNA of the Ti plasmid pTiACH 5 (Gielen et al., EMBO J. 3 (1984), 835-846), nucleotides 11749-11939, which had been isolated as a PvuII/HindIII fragment from the plasmid pAGV 40 (Herrera-Estrella et al., Nature 303 (1983), 209-213) and which after addition of SphI linkers to the PvuII cleavage site had been cloned between the SphI and HindIII cleavage site of the polylinker of pBin19.

The plasmid pBinRoIC-CiSy has a length of about 13 kb.

The plasmid pBinRoIC-CiSy was inserted into potato plants via the gene transfer mediated by *Agrobacterium tumefaciens*. Whole plants were regenerated from transformed cells.

The analysis of a number of plants transformed with this vector system unambiguously showed an increased biomass, which is a result of the expression of the CiSy cDNA from yeast contained in pBinRoIC-CiSy.

### **Example 7**

#### **Grafting experiment**

For grafting the shoot of a receiver plant is replaced with the shoot of a donor plant.

In this experiment the shoot of a transgenic plant (RoIC-Suc2 #25) is grafted onto the stock of a wildtype plant (*Solanum tuberosum*, var. Désirée). In a control experiment a wildtype shoot is grafted onto a wildtype stock in order to rule out culturing differences in the experiments (autografting). The aim of the experiment is to examine the exclusive impact of the photosynthetic activity and photoassimilate distribution of a transgenic shoot on organs (in this case tubers) of a wildtype stock.

Potato plants were transferred from a tissue culture to soil and placed into a greenhouse. After approx. five weeks (the plants have not yet induced tuber production at this stage) the plants are grafted. For this purpose the shoot of the receiver plant which is not needed is cut off, and a wedge is cut into the stem of the receiver plant. The donor shoot to be grafted is cut at the stem end in the appropriate way and is inserted into the wedge of the receiver plant. The grafting site is fixed with an adhesive tape.

Then the grafted potato plants are kept under increased air humidity and in shadow for approx. one week. Within seven to ten days they are step by step adapted to normal greenhouse conditions. At this stage the plants are seven weeks old.

All leaves of the receiver plant are now removed, and the stem is covered from light with an aluminum sheet in order to guarantee that exclusively the photosynthetic activity and the photoassimilate distribution of the donor shoot nourishes the stock of the grafted plant.

The plants are kept in the greenhouse until the potato tubers are harvested approx. two months after the grafting and approx. three months after the planting into soil. The results of such a grafting experiment are illustrated in Figure 12.

## Sequence Listing

## (1) GENERAL INFORMATION:

## (i) APPLICANT:

- (A) NAME: Max-Planck-Gesellschaft zur Foerderung der  
Wissenschaften e.V
- (B) STREET: none
- (C) CITY: Berlin
- (E) COUNTRY: Germany
- (F) POSTAL CODE: none

(ii) TITLE OF INVENTION: Process for increasing the yield in plants

(iii) NUMBER OF SEQUENCES: 1

## (iv) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- (D) SOFTWARE: PatentIn Release #1.0, Version #1.30 (EPA)

## (2) INFORMATION FOR SEQ ID NO: 1:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1138 base pairs
- (B) TYPE: nucleotide
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Genomic DNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Agrobacterium rhizogenes

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

AATTCGATAC GAAAAAGGCA AGTGCCAGGG CCATTTAAAA TACGGCGTCG GAAACTGGCG	60
CCAATCAGAC ACAGTCTCTG GTCGGGAAAG CCAGAGGTAG TTTGGCAACA ATCACATCAA	120
GATCGATGCG CAAGACACGG GAGGCCTTAA AATCTGGATC AAGCGAAAAT ACTGCATGCG	180
TGATCGTTCA TGGGTTTCATA GTACTGGGTT TGCTTTTTCT TGTCGTGTTG TTTGGCCTTA	240
GCGAAAGGAT GTCAAAAAAG GATGCCCATA ATTGGGAGGA GTGGGGTAAA GCTTAAAGTT	300
GGCCCGCTAT TGGATTTTCG GAAAGCGGCA TTGGCAAACG TGAAGATTGC TGCATTCAAG	360
ATACTTTTTC TATTTTCTGG TTAAGATGTA AAGTATTGCC ACAATCATAT TAATTACTAA	420
CATTGTATAT GTAATATAGT GCGGAAATTA TCTATGCCAA AATGATGTAT TAATAATAGC	480

AATAATAATA	TGTGTTAATC	TTTTTCAATC	GGAATACGT	TTAAGCGATT	ATCGTGTTGA	540
ATAAATTATT	CCAAAAGGAA	ATACATGGTT	TTGGAGAACC	TGCTATAGAT	ATATGCCAAA	600
TTTACACTAG	TTTAGTGGGT	GCAAAACTAT	TATCTCTGTT	TCTGAGTTTA	ATAAAAAATA	660
AATAAGCAGG	GCGAATAGCA	GTTAGCCTAA	GAAGGAATGG	TGGCCATGTA	CGTGCTTTTA	720
AGAGACCCTA	TAATAAATTG	CCAGCTGTGT	TGCTTTGGTG	CCGACAGGCC	TAACGTGGGG	780
TTTAGCTTGA	CAAAGTAGCG	CCTTTCCGCA	GCATAAATAA	AGGTAGGCGG	GTGCGTCCCA	840
TTATTAAAGG	AAAAAGCAAA	AGCTGAGATT	CCATAGACCA	CAAACCACCA	TTATTGGAGG	900
ACAGAACCTA	TTCCCTCACG	TGGGTCGCTA	GCTTTAAACC	TAATAAGTAA	AAACAATTAA	960
AAGCAGGCAG	GTGTCCCTTC	TATATTGCGA	CAACGAGGCG	ACGTGGAGCA	TCGACAGCCG	1020
CATCCATTAA	TTAATAAATT	TGTGGACCTA	TACCTAACTC	AAATATTTTT	ATTATTTGCT	1080
CCAATACGCT	AAGAGCTCTG	GATTATAAAT	AGTTTGGATG	CTTCGAGTTA	TGGGGTAC	1138



### Claims

1. A process for increasing the yield of plants, characterized in that recombinant DNA molecules containing
  - (a) a region allowing the transcription specifically in the companion cells; and operatively linked thereto
  - (b) a nucleotide sequence encoding a polypeptide selected from the group consisting of:
    - (i) proteins with an enzymatic activity that cleaves sucrose;
    - (ii) sucrose transporters;
    - (iii) proteins the activity of which leads to the stimulation of the proton gradients located at the plasma membrane of plant cells; and
    - (iv) citrate synthases;and which are stably integrated into the genome of plants are expressed.
2. The process of claim 1, wherein the nucleotide sequence encodes a plant protein.
3. The process of claim 1, wherein the nucleotide sequence encodes a protein from a bacterium or a fungus.
4. The process of claim 1, wherein the nucleotide sequence encodes a protein with an enzymatic activity that cleaves sucrose, selected from the group consisting of sucrose synthases, sucrose phosphorylases and invertases.
5. The process of claim 1, wherein the nucleotide sequence encodes a sucrose transporter from *Spinacia oleracea*.
6. The process of claim 1, wherein the nucleotide sequence encodes a proton ATPase.
7. The process of claim 6, wherein the nucleotide sequence encodes a proton ATPase from *Solanum tuberosum* or from *Saccharomyces cerevisiae*.


8. The process of any one of claims 1 to 7, wherein the region mentioned in (a) is the rolC promoter from *Agrobacterium rhizogenes*.
9. A recombinant nucleic acid molecule containing
  - (a) a region allowing the transcription specifically in the companion cells of plants; and operatively linked thereto
  - (b) a nucleotide sequence encoding a polypeptide, selected from the group consisting of
    - (i) sucrose synthases;
    - (ii) sucrose phosphorylases;
    - (iii) sucrose transporters;
    - (iv) proteins the activity of which leads to the stimulation of the proton gradient located at the plasma membrane of plant cells; and
    - (v) citrate synthases.
10. A vector containing a recombinant nucleic acid molecule of claim 9.
11. The vector of claim 10 which is suitable for the transformation of plant cells and for integration of foreign DNA into the plant genome.
12. A plant cell transformed with and containing a recombinant nucleic acid molecule of claim 9.
13. A plant containing plant cells of claim 12, wherein the plant shows an increased yield in comparison to a corresponding non-transformed plant due to the expression of the recombinant nucleic acid molecule in the companion cells of the plant.
14. Propagation material of a plant of claim 13 containing plant cells of claim 12.
15. Use of a recombinant nucleic acid molecule containing a region allowing the transcription specifically in the companion cells of plants and operatively linked

thereto a nucleotide sequence encoding a polypeptide selected from the group consisting of

- (i) proteins with an enzymatic activity that cleaves sucrose;
- (ii) sucrose transporters;
- (iii) proteins the activity of which leads to the stimulation of the proton gradient located at the plasma membrane; and
- (iv) citrate synthases,

for the expression in transgenic plants for increasing the yield.

1/17

**Figure 1**

2/17






Figure 2a


3/17

**Figure 2b**

4/17

**Figure 2c**


5/17




**Figure 3**



6/17

**Figure 4**

7/17

**Figure 5**

8/17

M #5 #7 #18 #19 #20 WT






Figure 6

9/17



**Figure 7**

10/17

**Figure 8**


11/17


#3 #5 #9 #11 #26 WT M



**Figure 9**

12/17

**Figure 10**



**Figure 11**








Figure 12

**Figure 13**

**Figure 14**

17/17




Figure 15

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 98/04878

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/82 C12N9/10 C12N9/24 C12N9/88 C12N9/14  
C12N15/54 C12N15/55 C12N15/60 C12N5/10 A01H5/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N A01H

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>LERCHL, J., ET AL. : "IMPAIRED PHOTOASSIMILATE PARTITIONING CAUSED BY PHLOEM-SPECIFIC REMOVAL OF PYROPHOSPHATE CAN BE COMPLEMENTED BY A PHLOEM-SPECIFIC CYTOSOLIC YEAST-DERIVED INVERTASE IN TRANSGENIC PLANTS"</p> <p>THE PLANT CELL, vol. 7, March 1995, pages 259-270, XP002086882</p> <p>especially page 259; Fig. 2 + 3</p> <p>see the whole document</p> <p style="text-align: center;">---</p> <p style="text-align: center;">-/--</p>	1, 3, 4, 8, 15

☒ Further documents are listed in the continuation of box C.

☐ Patent family members are listed in annex.

### \* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

7 December 1998

Date of mailing of the international search report

21/12/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl.  
Fax: (+31-70) 340-3016

Authorized officer

Holtorf, S

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 98/04878

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No
A	ZRENNER, R., ET AL. : "evidence of the crucial role of Sucrose synthase for sink strength using transgenic potato plants ( <i>Solanum tuberosum</i> L.)" THE PLANT JOURNAL, vol. 7, no. 1, 1995, pages 97-107, XP002086883 page 104, right column; Fig. 9 ---	1-15
A	RIESMEIER, J.W., ET AL. : "evidence for an essential role of the sucrose transpotrer in phloem loading and assimilate partitioning" THE EMBO JOURNAL, vol. 13, no. 1, 1994, pages 1-7, XP002086884 abstract, Fig. 2; Table 1; pages 5,6; page 6, last sentence ---	1-15
A	FOYER C H ET AL: "MODIFICATIONS IN CARBON ASSIMILATION, CARBON PARTITIONING AND TOTAL BIOMASS AS A RESULT OF OVER-EXPRESSION OF SUCROSE PHOSPHATES SYNTHASE IN TRANSGENIC TOMATO PLANTS" PLANT PHYSIOLOGY, vol. 105, no. 1, 1 May 1994, page 23 XP002002230 see the whole document ---	1-15
A	MARTIN T ET AL: "EXPRESSION OF AN ARABIDOPSIS SUCROSE SYNTHASE GENE INDICATES A ROLE IN METABOLIZATION OF SUCROSE BOTH DURING PHLOEM LOADING AND IN SINK ORGANS" PLANT JOURNAL, vol. 4, no. 2, 1993, pages 367-377, XP002061421 abstract, page 307, right column; page 368, left column; page 371,373,375; Fig. 7 ---	1-15
A	SONNEWALD. U., ET AL. : "increased potato tuber size resulting from apoplastic expression of a yeast invertase" NATURE BIOTECHNOLOGY, vol. 15, August 1997, pages 794-797, XP002086885 see the whole document ---	1-15
	--- -/--	

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 98/04878

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>RIESMEIER, J.W., ET AL. : "potato sucrose transporter expression in minor veins indicates a role in phloem loading"</p> <p>THE PLANT CELL,</p> <p>vol. 5, November 1993, pages 1591-1598,</p> <p>XP002086886</p> <p>see the whole document</p> <p>-----</p>	1-15